

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 April 2003 (03.04.2003)

PCT

(10) International Publication Number
WO 03/026631 A1

(51) International Patent Classification⁷: A61K 9/72,
31/192, 31/407, 31/405, 31/473

(21) International Application Number: PCT/US02/18543

(22) International Filing Date: 13 May 2002 (13.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/294,203 24 May 2001 (24.05.2001) US
60/317,479 5 September 2001 (05.09.2001) US

(71) Applicant: ALEXZA MOLECULAR DELIVERY
CORPORATION [US/US]; 1001 E.Meadow Circle, Palo
Alto, CA 94303 (US).

(72) Inventors: RABINOWITZ, Joshua, D.; 750 N. Shore-
line Blvd. #50, Mountain View, CA 94043 (US). ZAF-
FARONI, Alejandro, C.; 1 Faxon Forest, Atherton, CA
94027 (US).

(74) Agents: ECKMAN, Richard, R. et al.; Morrison & Foer-
ster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018
(US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: DELIVERY OF DRUG ESTERS THROUGH AN INHALATION ROUTE

(57) Abstract: The present invention relates to the delivery of drug esters through an inhalation route. Specifically, it relates to aerosols containing drug esters that are used in inhalation therapy. In a composition aspect of the present invention, the aerosol comprises particles comprising at least 5 percent by weight of drug ester. In a method aspect of the present invention, a drug ester is delivered to a mammal through an inhalation route. The method comprises: a) heating a composition, wherein the composition comprises at least 5 percent by weight of drug ester, to form a vapor; and, b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal. In a kit aspect of the present invention, a kit for delivering a drug ester through an inhalation route to a mammal is provided which comprises: a) a composition comprising at least 5 percent by weight of drug ester; and, b) a device that forms a drug ester containing aerosol from the composition, for inhalation by the mammal.

WO 03/026631 A1

DELIVERY OF DRUG ESTERS THROUGH AN INHALATION ROUTE

This application claims priority to U.S. provisional application Ser. No. 60/294,203 entitled "Thermal Vapor Delivery of Drugs," filed May 24, 2001 and to U.S. provisional application Ser. No. 60/317,479 entitled "Aerosol Drug Delivery," filed September 5, 2001, both of which are hereby incorporated by reference for all purposes.

Field of the Invention

The present invention relates to the delivery of drug esters through an inhalation route. Specifically, it relates to aerosols containing drug esters that are used in inhalation therapy.

Background of the Invention

There are a number of compounds containing acids and alcohols that are currently marketed as drugs. In certain circumstances, the presence of such functionality prevents effective drug delivery. This phenomenon could be due to a range of effects, including poor solubility and inadequate transcellular transport.

It is desirable to provide a new route of administration for drug acids and alcohols that rapidly produces peak plasma concentrations of the compounds. The provision of such a route is an object of the present invention.

Summary of the Invention

The present invention relates to the delivery of drug esters through an inhalation route. Specifically, it relates to aerosols containing drug esters that are used in inhalation therapy.

In a composition aspect of the present invention, the aerosol comprises particles comprising at least 5 percent by weight of drug ester. Preferably, the drug ester has a decomposition index less than 0.15. More preferably, it has a decomposition index less than 0.10 or 0.05. Preferably, the particles comprise at least 10 percent by weight of drug ester. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent or 99.97 percent by weight of drug ester.

Typically, the drug ester is an ester of a drug from one of the following classes: antibiotics, anticonvulsants, antidepressants, antihistamines, antiparkinsonian drugs, drugs for migraine headaches, drugs for the treatment of alcoholism, muscle relaxants, anxiolytics, nonsteroidal anti-inflammatories, other analgesics and steroids.

Typically, where the drug ester is an ester of an antibiotic, it is selected from an ester of one of the following compounds: cefmetazole; cefazolin; cephalixin; cefoxitin; cephradine; cephaloglycin; cephaloridine; cephalosporins, such as cephalosporin c; cephalotin; cephamycins, such as cephamycin a, cephamycin b, and cephamycin c; cepharin; cephradine; ampicillin; amoxicillin; hetacillin; carfecillin; carindacillin; carbenicillin; amylpenicillin; azidocillin; benzylpenicillin; clometocillin; cloxacillin; cyclacillin; methicillin; nafcillin; 2-pentenylpenicillin; penicillins, such as penicillin n, penicillin o, penicillin s, and penicillin v; chlorobutin penicillin; dicloxacillin; diphenicillin; heptylpenicillin; and metampicillin.

Typically, where the drug ester is an ester of an anticonvulsant, it is selected from an ester of one of the following compounds: 4-amino-3-hydroxybutyric acid, ethanedisulfonate, gabapentin, and vigabatrin.

Typically, where the drug ester is an ester of an antidepressant, it is selected from an ester of one of the following compounds: tianeptine and S-adenosylmethionine.

Typically, where the drug ester is an ester of an antihistamine, it is an ester of fexofenadine.

Typically, where the drug ester is an ester of an antiparkinsonian drug, it is selected from an ester of one of the following compounds: apomorphine, baclofen, levodopa, carbidopa, and thioctate.

Typically, where the drug ester is an ester of a drug for migraine headaches, it is selected from an ester of one of the following compounds: aspirin, diclofenac, naproxen, tolfenamic acid, and valproate.

Typically, where the drug ester is an ester of a drug for the treatment of alcoholism, it is an ester of acamprosate.

Typically, where the drug ester is an ester of a muscle relaxant, it is an ester of baclofen.

Typically, where the drug ester is an ester of an anxiolytic, it is selected from an ester of one of the following compounds: chlorazepate, calcium N-carboamoylaspartate and chloral betaine.

Typically, where the drug ester is an ester of a nonsteroidal anti-inflammatory, it is selected from an ester of one of the following compounds: aceclofenac, alclofenac, alminoprofen, amfenac, aspirin, benoxaprofen, bermoprofen, bromfenac, bufexamac, butibufen, bucloxate, carprofen, cinchophen, cinmetacin, clidanac, clopriac, clometacin, diclofenac, diflunisal, etodolac, fenclozate, fenoprofen, flutiazin, flurbiprofen, ibuprofen, ibufenac, indomethacin, indoprofen, ketoprofen, ketorolac, loxoprofen, meclofenamate, naproxen, oxaprozin, pirprofen, prodolic acid, salsalate, sulindac, tofenamate, and tolmetin.

Typically, where the drug ester is an ester of an other analgesic, it is selected from an ester of one of the following compounds: bumadizon, clometacin, and clonixin.

Typically, where the drug ester is an ester of a steroid, it is selected from an ester of one of the following compounds: betamethasone, chloroprednisone, clocortolone, cortisone, desonide, dexamethasone, desoximetasone, difluprednate, estradiol, fludrocortisone, flumethasone, flunisolide, fluocortolone, fluprednisolone, hydrocortisone, meprednisone, methylprednisolone, paramethasone, prednisolone, prednisone, pregnan-3-alpha-ol-20-one, testosterone, and triamcinolone.

Typically, where the drug ester is an ester of a drug acid, the ester is selected from an ester of the following type: C₁-C₆ straight chain substituted or unsubstituted alkyl ester, C₁-C₆ branched chain substituted or unsubstituted alkyl ester, C₃-C₆ substituted or unsubstituted cyclic alkyl ester, C₁-C₆ substituted or unsubstituted alkenyl ester, C₁-C₆ substituted or unsubstituted alkynyl ester, and substituted or unsubstituted aromatic ester.

Typically, where the drug ester is an ester of a drug alcohol, the ester is selected from an ester of the following type: C₁-C₆ substituted or unsubstituted straight chain alkanoate, C₁-C₆ substituted or unsubstituted branched chain alkanoate, C₁-C₆ substituted or unsubstituted alkenoate, and C₁-C₆ substituted or unsubstituted alkynoate.

Typically, the drug ester is selected from one of the following: ketoprofen methyl ester, ketoprofen ethyl ester, ketoprofen norcholine ester, ketorolac methyl ester, ketorolac ethyl ester, ketorolac norcholine ester, indomethacin methyl ester, indomethacin ethyl ester, indomethacine norcholine ester, and apomorphine diacetate.

Typically, the aerosol has a mass of at least 0.01 mg. Preferably, the aerosol has a mass of at least 0.05 mg. More preferably, the aerosol has a mass of at least 0.10 mg, 0.15 mg, 0.2 g or 0.25 mg.

Typically, the particles comprise less than 10 percent by weight of drug ester degradation products. Preferably, the particles comprise less than 5 percent by weight of

drug ester degradation products. More preferably, the particles comprise less than 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of drug ester degradation products.

Typically, the particles comprise less than 90 percent by weight of water.

Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

Typically, the aerosol has an inhalable aerosol drug ester mass density of between 0.1 mg/L and 100 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 0.1 mg/L and 75 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 0.1 mg/L and 50 mg/L.

Typically, the aerosol has an inhalable aerosol particle density greater than 10^6 particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than 10^7 particles/mL or 10^8 particles/mL.

Typically, the aerosol particles have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 2. Preferably, the geometric standard deviation is less than 1.9. More preferably, the geometric standard deviation is less than 1.8, 1.7, 1.6 or 1.5.

Typically, the aerosol is formed by heating a composition containing drug ester to form a vapor and subsequently allowing the vapor to condense into an aerosol.

In a method aspect of the present invention, a drug ester is delivered to a mammal through an inhalation route. The method comprises: a) heating a composition, wherein the composition comprises at least 5 percent by weight of drug ester, to form a vapor; and, b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal. Preferably, the drug ester has a decomposition index less than 0.15. More preferably, it has a decomposition index less than 0.10 or 0.05.

Preferably, the composition that is heated comprises at least 10 percent by weight of drug ester. More preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of drug ester.

Typically, the drug ester is an ester of a drug from one of the following classes: antibiotics, anticonvulsants, antidepressants, antihistamines, antiparkinsonian drugs, drugs for migraine headaches, drugs for the treatment of alcoholism, muscle relaxants, anxiolytics, nonsteroidal anti-inflammatories, other analgesics and steroids.

5 Typically, where the drug ester is an ester of an antibiotic, it is selected from an ester of one of the following compounds: cefmetazole; cefazolin; cephalixin; cefoxitin; cephacetrile; cephaloglycin; cephaloridine; cephalosporins, such as cephalosporin c; cephalotin; cephamycins, such as cephamycin a, cephamycin b, and cephamycin c; cepharin; cephradine; ampicillin; amoxicillin; hetacillin; carfecillin; carindacillin;
10 carbenicillin; amylpenicillin; azidocillin; benzylpenicillin; clometocillin; cloxacillin; cyclacillin; methicillin; nafcillin; 2-pentenylpenicillin; penicillins, such as penicillin n, penicillin o, penicillin s, and penicillin v; chlorobutin penicillin; dicloxacillin; diphenicillin; heptylpenicillin; and metampicillin.

15 Typically, where the drug ester is an ester of an anticonvulsant, it is selected from an ester of one of the following compounds: 4-amino-3-hydroxybutyric acid, ethanedisulfonate, gabapentin, and vigabatrin.

Typically, where the drug ester is an ester of an antidepressant, it is selected from an ester of one of the following compounds: tianeptine and S-adenosylmethionine.

20 Typically, where the drug ester is an ester of an antihistamine, it is an ester of fexofenadine.

Typically, where the drug ester is an ester of an antiparkinsonian drug, it is selected from an ester of one of the following compounds: apomorphine, baclofen, levodopa, carbidopa, and thioctate.

25 Typically, where the drug ester is an ester of a drug for migraine headaches, it is selected from an ester of one of the following compounds: aspirin, diclofenac, naproxen, tolfenamic acid, and valproate.

Typically, where the drug ester is an ester of a drug for the treatment of alcoholism, it is an ester of acamprosate.

30 Typically, where the drug ester is an ester of a muscle relaxant, it is an ester of baclofen.

Typically, where the drug ester is an ester of an anxiolytic, it is selected from an ester of one of the following compounds: chlorazepate, calcium N-carboamoylaspartate and chloral betaine.

Typically, where the drug ester is an ester of a nonsteroidal anti-inflammatory, it is selected from an ester of one of the following compounds: aceclofenac, alclofenac, alminoprofen, amfenac, aspirin, benoxaprofen, bermoprofen, bromfenac, bufexamac, butibufen, bucloxate, carprofen, cinchophen, cinmetacin, clidanac, clopiac, clometacin, diclofenac, diflunisal, etodolac, fenclozate, fenoprofen, flutiazin, flurbiprofen, ibuprofen, ibufenac, indomethacin, indoprofen, ketoprofen, ketorolac, loxoprofen, meclofenamate, naproxen, oxaprozin, pirprofen, prodolic acid, salsalate, sulindac, tofenamate, and tolmetin.

Typically, where the drug ester is an ester of an other analgesic, it is selected from an ester of one of the following compounds: bumadizon, clometacin, and clonixin.

Typically, where the drug ester is an ester of a steroid, it is selected from an ester of one of the following compounds: betamethasone, chloroprednisone, clocortolone, cortisone, desonide, dexamethasone, desoximetasone, difluprednate, estradiol, fludrocortisone, flumethasone, flunisolide, fluocortolone, fluprednisolone, hydrocortisone, meprednisone, methylprednisolone, paramethasone, prednisolone, prednisone, pregnan-3-alpha-ol-20-one, testosterone, and triamcinolone.

Typically, where the drug ester is an ester of a drug acid, the ester is selected from an ester of the following type: C₁-C₆ straight chain substituted or unsubstituted alkyl ester, C₁-C₆ branched chain substituted or unsubstituted alkyl ester, C₃-C₆ substituted or unsubstituted cyclic alkyl ester, C₁-C₆ substituted or unsubstituted alkenyl ester, C₁-C₆ substituted or unsubstituted alkynyl ester, and substituted or unsubstituted aromatic ester.

Typically, where the drug ester is an ester of a drug alcohol, the ester is selected from an ester of the following type: C₁-C₆ substituted or unsubstituted straight chain alkanoate, C₁-C₆ substituted or unsubstituted branched chain alkanoate, C₁-C₆ substituted or unsubstituted alkenoate, and C₁-C₆ substituted or unsubstituted alkynoate.

Typically, the drug ester is selected from one of the following: ketoprofen methyl ester, ketoprofen ethyl ester, ketoprofen norcholine ester, ketorolac methyl ester, ketorolac ethyl ester, ketorolac norcholine ester, indomethacin methyl ester, indomethacin ethyl ester, indomethacine norcholine ester, and apomorphine diacetate.

Typically, the particles comprise at least 5 percent by weight of drug ester.

Preferably, the particles comprise at least 10 percent by weight of drug ester. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of drug ester.

Typically, the condensation aerosol has a mass of at least 0.01 mg. Preferably, the aerosol has a mass of at least 0.05 mg. More preferably, the aerosol has a mass of at least 0.10 mg, 0.15 mg, 0.2 g or 0.25 mg.

Typically, the particles comprise less than 10 percent by weight of drug ester degradation products. Preferably, the particles comprise less than 5 percent by weight of drug ester degradation products. More preferably, the particles comprise 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of drug ester degradation products.

Typically, the particles comprise less than 90 percent by weight of water. Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

Typically, the particles of the delivered condensation aerosol have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 2. Preferably, the geometric standard deviation is less than 1.9. More preferably, the geometric standard deviation is less than 1.8, 1.7, 1.6 or 1.5.

Typically, the delivered aerosol has an inhalable aerosol drug ester mass density of between 0.1 mg/L and 100 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 0.1 mg/L and 75 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 0.1 mg/L and 50 mg/L.

Typically, the delivered aerosol has an inhalable aerosol particle density greater than 10^6 particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than 10^7 particles/mL or 10^8 particles/mL.

Typically, the rate of inhalable aerosol particle formation of the delivered condensation aerosol is greater than 10^8 particles per second. Preferably, the aerosol is formed at a rate greater than 10^9 inhalable particles per second. More preferably, the aerosol is formed at a rate greater than 10^{10} inhalable particles per second.

Typically, the delivered condensation aerosol is formed at a rate greater than 0.5 mg/second. Preferably, the aerosol is formed at a rate greater than 0.75 mg/second. More

preferably, the aerosol is formed at a rate greater than 1 mg/second, 1.5 mg/second or 2 mg/second.

Typically, between 0.1 mg and 100 mg of drug ester are delivered to the mammal in a single inspiration. Preferably, between 0.1 mg and 75 mg of drug ester are delivered to the mammal in a single inspiration. More preferably, between 0.1 mg and 50 mg of drug ester are delivered in a single inspiration.

Typically, the delivered condensation aerosol results in a peak plasma concentration of drug acid or drug alcohol in the mammal in less than 1 h. Preferably, the peak plasma concentration is reached in less than 0.5 h. More preferably, the peak plasma concentration is reached in less than 0.2, 0.1, 0.05, 0.02 or 0.01 h.

In a kit aspect of the present invention, a kit for delivering a drug ester through an inhalation route to a mammal is provided which comprises: a) a composition comprising at least 5 percent by weight of drug ester; and, b) a device that forms a drug ester aerosol from the composition, for inhalation by the mammal. Preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of drug ester.

Typically the drug ester has a decomposition index less than 0.15. More preferably, it has a decomposition index less than 0.10 or 0.05.

Typically, the device contained in the kit comprises: a) an element for heating the drug ester composition to form a vapor; b) an element allowing the vapor to cool to form an aerosol; and, c) an element permitting the mammal to inhale the aerosol.

Brief Description of the Figure

Fig. 1 shows a cross-sectional view of a device used to deliver drug ester aerosols to a mammal through an inhalation route.

Detailed Description of the Invention

Definitions

"Aerodynamic diameter" of a given particle refers to the diameter of a spherical droplet with a density of 1 g/mL (the density of water) that has the same settling velocity as the given particle.

"Aerosol" refers to a suspension of solid or liquid particles in a gas.

“Aerosol drug ester mass density” refers to the mass of drug ester per unit volume of aerosol.

“Aerosol mass density” refers to the mass of particulate matter per unit volume of aerosol.

5 “Aerosol particle density” refers to the number of particles per unit volume of aerosol.

“Condensation aerosol” refers to an aerosol formed by vaporization of a substance followed by condensation of the substance into an aerosol.

10 “Decomposition index” refers to a number derived from an assay described in Example 8. The number is determined by subtracting the percent purity of the generated aerosol from 1.

“Drug” refers to any chemical compound that is used in the prevention, diagnosis, treatment, or cure of disease, for the relief of pain, or to control or improve any physiological or pathological disorder in humans or animals. Such compounds are
15 oftentimes listed in the Physician’s Desk Reference (Medical Economics Company, Inc. at Montvale, NJ, 56th edition, 2002), which is herein incorporated by reference.

“Drug acid” refers to a drug containing a carboxylic acid moiety.

“Drug alcohol” refers to a drug containing a hydroxyl moiety.

20 “Drug Ester” refers to a drug acid or drug alcohol, where the carboxylic acid group or hydroxyl group has been chemically modified to form an ester. The drug acids and alcohols from which the esters are formed come from a variety of drug classes, including, without limitation, antibiotics, anticonvulsants, antidepressants, antihistamines, antiparkinsonian drugs, drugs for migraine headaches, drugs for the treatment of alcoholism, muscle relaxants, anxiolytics, nonsteroidal anti-inflammatories, other
25 analgesics, and steroids.

Examples of antibiotics from which drug esters are formed include cefmetazole; cefazolin; cephalixin; cefoxitin; cephacetrile; cephaloglycin; cephaloridine; cephalosporins, such as cephalosporin c; cephalotin; cephamycins, such as cephamycin a, cephamycin b, and cephamycin c; cepharin; cephradine; ampicillin; amoxicillin; hetacillin;
30 carfecillin; carindacillin; carbenicillin; amylpenicillin; azidocillin; benzylpenicillin; clometocillin; cloxacillin; cyclacillin; methicillin; nafcillin; 2-pentenylpenicillin; penicillins, such as penicillin n, penicillin o, penicillin s, and penicillin v; chlorobutin penicillin; dicloxacillin; diphenicillin; heptylpenicillin; and metampicillin.

Examples of anticonvulsants from which drug esters are formed include 4-amino-3-hydroxybutyric acid, ethanedisulfonate, gabapentin, and vigabatrin.

Examples of antidepressants from which drug esters are formed include tianeptine and S-adenosylmethionine.

5 Examples of antihistamines from which drug esters are formed include fexofenadine.

Examples of antiparkinsonian drugs from which drug esters are formed include apomorphine, baclofen, levodopa, carbidopa, and thioclate.

10 Examples of anxiolytics from which drug esters are formed include chlorazepate, calcium N-carboamoylaspartate and chloral betaine.

Examples of drugs for migraine headache from which drug esters are formed include aspirin, diclofenac, naproxen, tolfenamic acid, and valproate.

Examples of drugs for the treatment of alcoholism from which drug esters are formed include acamprosate.

15 Examples of muscle relaxants from which drug esters are formed include baclofen.

Examples of nonsteroidal anti-inflammatories from which drug esters are formed include aceclofenac, alclofenac, alminoprofen, amfenac, aspirin, benoxaprofen, bermoprofen, bromfenac, bufexamac, butibufen, bucloxate, carprofen, cinchophen, cinmetacin, clidanac, clopiac, clometacin, diclofenac, diflunisal, etodolac, fenclozate, fenoprofen, flutiazin, flurbiprofen, ibuprofen, ibufenac, indomethacin, indoprofen, ketoprofen, ketorolac, loxoprofen, meclofenamate, naproxen, oxaprozin, pirprofen, prodolic acid, salsalate, sulindac, tofenamate, and tolmetin.

20

Examples of other analgesics from which drug esters are formed include bumadizon, clometacin, and clonixin.

25 Examples of steroids from which drug esters are formed include betamethasone, chloroprednisone, clocortolone, cortisone, desonide, dexamethasone, desoximetasone, difluprednate, estradiol, fludrocortisone, flumethasone, flunisolide, fluocortolone, fluprednisolone, hydrocortisone, meprednisone, methylprednisolone, paramethasone, prednisolone, prednisone, pregnan-3-alpha-ol-20-one, testosterone, and triamcinolone.

30 Examples of drug esters formed from drug acids include C₁-C₆ straight chain substituted or unsubstituted alkyl esters, C₁-C₆ branched chain substituted or unsubstituted alkyl esters, C₃-C₆ substituted or unsubstituted cyclic alkyl esters, C₁-C₆ substituted or unsubstituted alkenyl esters, C₁-C₆ substituted or unsubstituted alkynyl esters, and

substituted or unsubstituted aromatic esters. C₁-C₆ straight chain unsubstituted alkyl esters include, for example, methyl ester, ethyl ester and propyl ester. C₁-C₆ straight chain substituted alkyl esters include, for example, 2-(dimethylamino)-ethyl ester (-CH₂CH₂N(CH₃)₂). C₁-C₆ branched chain unsubstituted alkyl esters include, for example, isopropyl ester and isobutyl ester. C₁-C₆ branched chain substituted alkyl esters include, for example, 2-(dimethylamino)-isopropyl ester (-CH(CH₃)CH₂N(CH₃)₂). C₃-C₆ unsubstituted cyclic alkyl esters include, for example, cyclopropyl and cyclohexyl ester. C₃-C₆ substituted cyclic alkyl esters include, for example, 2-(dimethylamino)-cyclopropyl ester. C₁-C₆ unsubstituted alkenyl esters include, for example, 2-butenyl ester (-CH₂CHCHCH₃). C₁-C₆ substituted alkenyl esters include, for example, 4-(dimethylamino)-2-butenyl ester (-CH₂CHCHCH₂N(CH₃)₂). C₁-C₆ unsubstituted alkynyl esters include, for example, 2-butyne ester (-CH₂CCCH₃). C₁-C₆ substituted alkynyl esters include, for example, 4-(dimethylamino)-2-butyne ester (-CH₂CCCH₂N(CH₃)₂). Unsubstituted aromatic esters include, for example, phenyl ester and naphthyl ester. Substituted aromatic esters include, for example, 4-(dimethylamino)phenyl ester.

Examples of drug esters formed from drug alcohols include C₁-C₆ substituted or unsubstituted straight chain alkanoates, C₁-C₆ substituted or unsubstituted branched chain alkanoates, C₁-C₆ substituted or unsubstituted alkenoates, and C₁-C₆ substituted or unsubstituted alkynoates. C₁-C₆ unsubstituted straight chain alkanoates include, for example, methanoate (-C(O)H), ethanoate (-C(O)CH₃) and propanoate (-C(O)CH₂CH₃). C₁-C₆ substituted straight chain alkanoates include, for example, 2-(phenyl)-ethanoate (-C(O)CH₂Ph). C₁-C₆ unsubstituted branched chain alkanoates include, for example, isobutanoate (-C(O)CH(CH₃)₂). C₁-C₆ substituted branched chain alkanoates include, for example, 3-(phenyl)-isobutanoate (-C(O)CH(CH₃)CH₂Ph). C₁-C₆ unsubstituted alkenoates include, for example, 2-butenate (-C(O)CHCHCH₃). C₁-C₆ substituted alkenoates include, for example, 4-(phenyl)-2-butenate (-C(O)CHCHCH₂Ph). C₁-C₆ unsubstituted alkynoates include, for example, 2-butyneate (-C(O)CCCH₃). C₁-C₆ substituted alkynoates include, for example, 4-(phenyl)-2-butyneate.

Examples of other drug esters are found in U.S. Pat. No. 5,607,691 to Hale *et al.* and U.S. Pat. No. 5,622,944 to Hale *et al.* These patents are herein incorporated by reference.

"Drug ester degradation product" refers to a compound resulting from a chemical modification of the drug ester. The modification, for example, can be the result of a

thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

“Inhalable aerosol drug ester mass density” refers to the aerosol drug ester mass density produced by an inhalation device and delivered into a typical patient tidal volume.

5 “Inhalable aerosol mass density” refers to the aerosol mass density produced by an inhalation device and delivered into a typical patient tidal volume.

“Inhalable aerosol particle density” refers to the aerosol particle density of particles of size between 100 nm and 5 microns produced by an inhalation device and delivered into a typical patient tidal volume.

10 “Mass median aerodynamic diameter” or “MMAD” of an aerosol refers to the aerodynamic diameter for which half the particulate mass of the aerosol is contributed by particles with an aerodynamic diameter larger than the MMAD and half by particles with an aerodynamic diameter smaller than the MMAD.

15 “Norcholine ester” refers to an ester where the portion attached to the ester oxygen is $-\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$.

“Rate of aerosol formation” refers to the mass of aerosolized particulate matter produced by an inhalation device per unit time.

“Rate of inhalable aerosol particle formation” refers to the number of particles of size between 100 nm and 5 microns produced by an inhalation device per unit time.

20 “Rate of drug ester aerosol formation” refers to the mass of aerosolized, drug ester produced by an inhalation device per unit time.

“Settling velocity” refers to the terminal velocity of an aerosol particle undergoing gravitational settling in air.

25 “Substituted” alkyl, alkenyl, alkynyl or aryl refers to the replacement of one or more hydrogen atoms on the moiety (e.g., alkyl) with another group. Such groups include, without limitation, the following: halo, amino, alkylamino, dialkylamino, hydroxyl, cyano, nitro and phenyl.

“Typical patient tidal volume” refers to 1 L for an adult patient and 15 mL/kg for a pediatric patient.

30 “Vapor” refers to a gas, and “vapor phase” refers to a gas phase. The term “thermal vapor” refers to a vapor phase, aerosol, or mixture of aerosol-vapor phases, formed preferably by heating.

Formation of Drug Esters from Drug Acids or Drug Alcohols

Formation of drug esters from drug acids is typically accomplished by reacting the acid, or an activated derivative (e.g., acid chloride or mixed anhydride) with an appropriate alcohol under conditions well known to those of skill in the art. *See, for example*,
5 Streitweiser, A., Jr. and Heathcock, C. H. (1981) Introduction to Organic Chemistry, Macmillan Publishing Col., Inc., New York. Conversely, formation of drug esters from drug alcohols is typically accomplished by reacting the alcohol with an appropriate activated acid derivative (e.g., ClC(O)CH_3). *See Id.*

Formation of Drug Ester Containing Aerosols

Any suitable method is used to form the aerosols of the present invention. A preferred method, however, involves heating a composition comprising a drug ester to form a vapor, followed by cooling of the vapor such that it condenses to provide a drug ester comprising aerosol (condensation aerosol). The composition is heated in one of two forms:
15 as pure active compound (i.e., pure drug ester); or, as a mixture of active compound and a pharmaceutically acceptable excipient.

Pharmaceutically acceptable excipients may be volatile or nonvolatile. Volatile excipients, when heated, are concurrently volatilized, aerosolized and inhaled with drug ester. Classes of such excipients are known in the art and include, without limitation,
20 gaseous, supercritical fluid, liquid and solid solvents. The following is a list of exemplary carriers within the classes: water; terpenes, such as menthol; alcohols, such as ethanol, propylene glycol, glycerol and other similar alcohols; dimethylformamide; dimethylacetamide; wax; supercritical carbon dioxide; dry ice; and mixtures thereof.

Solid supports on which the composition is heated are of a variety of shapes.
25 Examples of such shapes include, without limitation, cylinders of less than 1.0 mm in diameter, boxes of less than 1.0 mm thickness and virtually any shape permeated by small (e.g., less than 1.0 mm-sized) pores. Preferably, solid supports provide a large surface to volume ratio (e.g., greater than 100 per meter) and a large surface to mass ratio (e.g., greater than 1 cm^2 per gram).

30 A solid support of one shape can also be transformed into another shape with different properties. For example, a flat sheet of 0.25 mm thickness has a surface to volume ratio of approximately 8,000 per meter. Rolling the sheet into a hollow cylinder of

1 cm diameter produces a support that retains the high surface to mass ratio of the original sheet but has a lower surface to volume ratio (about 400 per meter).

A number of different materials are used to construct the solid supports. Classes of such materials include, without limitation, metals, inorganic materials, carbonaceous materials and polymers. The following are examples of the material classes: aluminum, silver, gold, stainless steel, copper and tungsten; silica, glass, silicon and alumina; graphite, porous carbons, carbon yarns and carbon felts; polytetrafluoroethylene and polyethylene glycol. Combinations of materials and coated variants of materials are used as well.

Where aluminum is used as a solid support, aluminum foil is a suitable material. Examples of silica, alumina and silicon based materials include amorphous silica S-5631 (Sigma, St. Louis, MO), BCR171 (an alumina of defined surface area greater than 2 m²/g from Aldrich, St. Louis, MO) and a silicon wafer as used in the semiconductor industry. Carbon yarns and felts are available from American Kynol, Inc., New York, NY. Chromatography resins such as octadecyl silane chemically bonded to porous silica are exemplary coated variants of silica.

The heating of the drug ester compositions is performed using any suitable method. Examples of methods by which heat can be generated include the following: passage of current through an electrical resistance element; absorption of electromagnetic radiation, such as microwave or laser light; and, exothermic chemical reactions, such as exothermic solvation, hydration of pyrophoric materials and oxidation of combustible materials.

Delivery of Drug Ester Containing Aerosols

Drug ester containing aerosols of the present invention are delivered to a mammal using an inhalation device. Where the aerosol is a condensation aerosol, the device has at least three elements: an element for heating a drug ester containing composition to form a vapor; an element allowing the vapor to cool, thereby providing a condensation aerosol; and, an element permitting the mammal to inhale the aerosol. Various suitable heating methods are described above. The element that allows cooling is, in its simplest form, an inert passageway linking the heating means to the inhalation means. The element permitting inhalation is an aerosol exit portal that forms a connection between the cooling element and the mammal's respiratory system.

One device used to deliver the drug ester containing aerosol is described in reference to Fig. 1. Delivery device 100 has a proximal end 102 and a distal end 104, a

heating module 106, a power source 108, and a mouthpiece 110. A drug ester composition is deposited on a surface 112 of heating module 106. Upon activation of a user activated switch 114, power source 108 initiates heating of heating module 106 (*e.g.*, through ignition of combustible fuel or passage of current through a resistive heating element). The drug ester composition volatilizes due to the heating of heating module 106 and condenses to form a condensation aerosol prior to reaching the mouthpiece 110 at the proximal end of the device 102. Air flow traveling from the device distal end 104 to the mouthpiece 110 carries the condensation aerosol to the mouthpiece 110, where it is inhaled by the mammal.

Devices, if desired, contain a variety of components to facilitate the delivery of drug ester containing aerosols. For instance, the device may include any component known in the art to control the timing of drug aerosolization relative to inhalation (*e.g.*, breath-actuation), to provide feedback to patients on the rate and/or volume of inhalation, to prevent excessive use (*i.e.*, "lock-out" feature), to prevent use by unauthorized individuals, and/or to record dosing histories.

In Vivo Hydrolysis of Drug Esters

After delivery of a drug ester aerosol to the lung of an animal, the ester moiety is typically hydrolyzed to provide the corresponding drug acid or drug alcohol, which produces a desired therapeutic effect. Where the ester reacts with water at \sim pH 7.4 at an appreciable rate, hydrolysis is chemically mediated. For other esters, hydrolysis is enzymatically mediated through the action of enzymes endogenous to the animal.

Dosage of Drug Ester Containing Aerosols

A typical dosage of a drug ester aerosol is either administered as a single inhalation or as a series of inhalations taken within an hour or less (dosage equals sum of inhaled amounts). Where the drug ester is administered as a series of inhalations, a different amount may be delivered in each inhalation. The dosage amount of drug ester in aerosol form is generally no greater than twice the standard dose of the drug acid or drug alcohol given orally.

One can determine the appropriate dose of drug ester containing aerosols to treat a particular condition using methods such as animal experiments and a dose-finding (Phase I/II) clinical trial. One animal experiment involves measuring plasma concentrations of drug acid or drug alcohol in an animal after its exposure to the aerosol. Mammals such as

dogs or primates are typically used in such studies, since their respiratory systems are similar to that of a human. Initial dose levels for testing in humans is generally less than or equal to the dose in the mammal model that resulted in plasma drug levels associated with a therapeutic effect in humans. Dose escalation in humans is then performed, until either an optimal therapeutic response is obtained or a dose-limiting toxicity is encountered.

Analysis of Drug Ester Containing Aerosols

Purity of a drug ester containing aerosol is determined using a number of methods, examples of which are described in Sekine *et al.*, *Journal of Forensic Science* 32:1271-1280 (1987) and Martin *et al.*, *Journal of Analytic Toxicology* 13:158-162 (1989). One method involves forming the aerosol in a device through which a gas flow (*e.g.*, air flow) is maintained, generally at a rate between 0.4 and 60 L/min. The gas flow carries the aerosol into one or more traps. After isolation from the trap, the aerosol is subjected to an analytical technique, such as gas or liquid chromatography, that permits a determination of composition purity.

A variety of different traps are used for aerosol collection. The following list contains examples of such traps: filters; glass wool; impingers; solvent traps, such as dry ice-cooled ethanol, methanol, acetone and dichloromethane traps at various pH values; syringes that sample the aerosol; empty, low-pressure (*e.g.*, vacuum) containers into which the aerosol is drawn; and, empty containers that fully surround and enclose the aerosol generating device. Where a solid such as glass wool is used, it is typically extracted with a solvent such as ethanol. The solvent extract is subjected to analysis rather than the solid (*i.e.*, glass wool) itself. Where a syringe or container is used, the container is similarly extracted with a solvent.

The gas or liquid chromatograph discussed above contains a detection system (*i.e.*, detector). Such detection systems are well known in the art and include, for example, flame ionization, photon absorption and mass spectrometry detectors. An advantage of a mass spectrometry detector is that it can be used to determine the structure of drug ester degradation products.

Particle size distribution of a drug ester containing aerosol is determined using any suitable method in the art (*e.g.*, cascade impaction). An Andersen Eight Stage Non-viable Cascade Impactor (Andersen Instruments, Smyrna, GA) linked to a furnace tube by a mock

throat (USP throat, Andersen Instruments, Smyrna, GA) is one system used for cascade impaction studies.

Inhalable aerosol mass density is determined, for example, by delivering a drug-containing aerosol into a confined chamber via an inhalation device and measuring the mass collected in the chamber. Typically, the aerosol is drawn into the chamber by having a pressure gradient between the device and the chamber, wherein the chamber is at lower pressure than the device. The volume of the chamber should approximate the tidal volume of an inhaling patient.

Inhalable aerosol drug ester mass density is determined, for example, by delivering a drug ester-containing aerosol into a confined chamber via an inhalation device and measuring the amount of non-degraded drug ester collected in the chamber. Typically, the aerosol is drawn into the chamber by having a pressure gradient between the device and the chamber, wherein the chamber is at lower pressure than the device. The volume of the chamber should approximate the tidal volume of an inhaling patient. The amount of non-degraded drug ester collected in the chamber is determined by extracting the chamber, conducting chromatographic analysis of the extract and comparing the results of the chromatographic analysis to those of a standard containing known amounts of drug ester.

Inhalable aerosol particle density is determined, for example, by delivering aerosol phase drug ester into a confined chamber via an inhalation device and measuring the number of particles of given size collected in the chamber. The number of particles of a given size may be directly measured based on the light-scattering properties of the particles. Alternatively, the number of particles of a given size is determined by measuring the mass of particles within the given size range and calculating the number of particles based on the mass as follows: Total number of particles = Sum (from size range 1 to size range N) of number of particles in each size range. Number of particles in a given size range = Mass in the size range/Mass of a typical particle in the size range. Mass of a typical particle in a given size range = $\pi \cdot D^3 \cdot \phi / 6$, where D is a typical particle diameter in the size range (generally, the mean boundary MMADs defining the size range) in microns, ϕ is the particle density (in g/mL) and mass is given in units of picograms (g^{-12}).

Rate of inhalable aerosol particle formation is determined, for example, by delivering aerosol phase drug ester into a confined chamber via an inhalation device. The delivery is for a set period of time (e.g., 3 s), and the number of particles of a given size collected in the chamber is determined as outlined above. The rate of particle formation is

equal to the number of 100 nm to 5 micron particles collected divided by the duration of the collection time.

Rate of aerosol formation is determined, for example, by delivering aerosol phase drug ester into a confined chamber via an inhalation device. The delivery is for a set period of time (*e.g.*, 3 s), and the mass of particulate matter collected is determined by weighing the confined chamber before and after the delivery of the particulate matter. The rate of aerosol formation is equal to the increase in mass in the chamber divided by the duration of the collection time. Alternatively, where a change in mass of the delivery device or component thereof can only occur through release of the aerosol phase particulate matter, the mass of particulate matter may be equated with the mass lost from the device or component during the delivery of the aerosol. In this case, the rate of aerosol formation is equal to the decrease in mass of the device or component during the delivery event divided by the duration of the delivery event.

Rate of drug ester aerosol formation is determined, for example, by delivering a drug ester containing aerosol into a confined chamber via an inhalation device over a set period of time (*e.g.*, 3 s). Where the aerosol is pure drug ester, the amount of drug collected in the chamber is measured as described above. The rate of drug ester aerosol formation is equal to the amount of drug ester aerosol collected in the chamber divided by the duration of the collection time. Where the drug ester containing aerosol comprises a pharmaceutically acceptable excipient, multiplying the rate of aerosol formation by the percentage of drug ester in the aerosol provides the rate of drug aerosol formation.

Utility of Drug Ester Containing Aerosols

The drug ester containing aerosols of the present invention are typically used for the same indication as the corresponding drug acid or drug alcohol. For instance, a drug ester of baclofen would be used to treat parkinsons disease and a drug ester of fexofenadine would be used to treat allergy symptoms.

The following examples are meant to illustrate, rather than limit, the present invention.

Drug acids or drug alcohols are typically commercially available from Sigma (www.sigma-aldrich.com), obtained in tablet form from a pharmacy and extracted, or synthesized using well known methods in the art.

EXAMPLE 1*General Procedures for Esterifying a Drug Acid*

1) Drug acid (10 mmol) is dissolved in 90 mL of dichloromethane. To the solution is added 1 drop of dimethylformamide and 13 mmol of oxalyl chloride. The resulting mixture is allowed to stir 30 min. The mixture is concentrated to dryness on a rotary evaporator to provide a residue, to which 50 mL of an alcohol (e.g., methanol) is added. The alcoholic solution is concentrated to dryness to afford the desired drug ester.

2) Drug acid (6 mmol) is dissolved in 60 mL of dichloromethane. To the solution is added 1 drop of dimethylformamide and 9 mmol of oxalyl chloride. The resulting mixture is allowed to stir 1 h. The mixture is concentrated to dryness on a rotary evaporator to provide a residue, to which 47 mmol of an alcohol (e.g., $\text{HOCH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$) in 20 mL dichloromethane is added. The reaction mixture is diluted with 60 mL dichloromethane and subjected to a series of washings: 50 mL saturated aqueous NaCl followed by 50 mL saturated aqueous NaHCO_3 and 2X 50 mL saturated aqueous NaCl. The dichloromethane extract is dried over Na_2SO_4 , filtered, and concentrated on a rotary evaporator to provide the desired drug ester.

EXAMPLE 2*General Procedure for Esterifying a Drug Alcohol*

Drug alcohol (5 mmol) is dissolved in 50 mL of dichloromethane. To the solution is added 5.5 mmol Hünig's base and 10 mmol acetyl chloride. The reaction mixture is allowed to stir at room temperature for 1 hour. The mixture is washed with 50 mL saturated aqueous NaHCO_3 followed by 50 mL saturated aqueous NaCl. The dichloromethane extract is dried over Na_2SO_4 , filtered, and concentrated on a rotary evaporator to provide the desired drug ester.

EXAMPLE 3*Procedure for Diesterifying Apomorphine*

Apomorphine·HCl·½H₂O (300 mg) was suspended in 600 µL of acetic acid. The suspension was heated to 100 °C and then cooled to 50 °C. Acetyl chloride (1 mL) was added to the suspension, which was heated at 40 °C for 3 h. The reaction mixture was allowed to cool to room temperature. Dichloromethane (1-2 mL) was added and the mixture was allowed to stir overnight. The reaction mixture was diluted with dichloromethane, and the solvent was removed on a rotary evaporator. Toluene (10 mL) was added to the residue and subsequently removed on a rotary evaporator. The toluene addition/removal was repeated. The resulting solid residue was triturated with ether, providing 430 mg of a solid (mp 158-160 °C).

A portion of the solid (230 mg) was suspended in 50 mL of dichloromethane. The suspension was washed with saturated aqueous NaHCO₃. The dichloromethane layer was dried over Na₂SO₄, filtered and concentrated on a rotary evaporator to provide 190 mg of the desired free base (mp ~110 °C).

EXAMPLE 4*Procedure for Synthesis of 2-(N,N-Dimethylamino)Ethyl Ester of Ketorolac*

Ketorolac (255 mg), triethylamine (101 mg) and 2-(dimethylamino)ethanol (HOCH₂CH₂N(CH₃)₂, 380 mg) were added to 2 mL dichloromethane. The mixture was cooled to -25 °C to -20 °C for 15 min. BOP (464 mg) was added, and the reaction mixture was gradually allowed to warm to room temperature. See Kim, M. H. and Patel, D. V. (1994) Tet. Lett. 35: 5603-5606. The reaction mixture was diluted with 60 mL of dichloromethane and washed sequentially with saturated aqueous NaCl, saturated aqueous NaHCO₃ and then saturated aqueous NaCl. The dichloromethane extract was dried over Na₂SO₄, filtered, and concentrated on a rotary evaporator to provide 390 mg of the desired material.

EXAMPLE 5*General Procedure for Volatilizing Compounds from Halogen Bulb*

A solution of drug in approximately 120 μ L dichloromethane is coated on a 3.5 cm x 7.5 cm piece of aluminum foil (precleaned with acetone). The dichloromethane is allowed to evaporate. The coated foil is wrapped around a 300 watt halogen tube (Feit Electric Company, Pico Rivera, CA), which is inserted into a glass tube sealed at one end with a rubber stopper. Running 60 V of alternating current (driven by line power controlled by a variac) through the bulb for 5-12 s or 90 V for 2.5-3.5 s affords thermal vapor (including aerosol), which is collected on the glass tube walls. (When desired, the system is flushed through with argon prior to volatilization.) Reverse-phase HPLC analysis with detection by absorption of 225 nm light is used to determine the purity of the aerosol.

Table 1, which follows, provides data from drugs volatilized using the above-recited general procedure.

Table 1

COMPOUND	AEROSOL PURITY	AEROSOL MASS
Indomethacin Methyl Ester	99%	1.44 mg
Indomethacin Ethyl Ester	>99%	3.09 mg
Indomethacin Norcholine Ester	100%	2.94 mg
Ketoprofen Methyl Ester	99%	4.4 mg
Ketoprofen Ethyl Ester	99.65%	4.11 mg
Ketoprofen Norcholine Ester	100%	2.6 mg
Ketorolac Methyl Ester	100%	3.17 mg
Ketorolac Ethyl Ester	>99%	5.19 mg
Ketorolac Norcholine Ester	100%	1.64 mg
Apomorphine Diacetate-HCl	94%	1.65 mg
Apomorphine Diacetate	96.9%	2.03 mg

EXAMPLE 6*General Procedure for Hydrolysis Studies of Drug Esters*

Drug ester (20 μ L, 10 mM acetonitrile) is added to 1 mL PBS solution (pH 7.5) at room temperature. At intermittent time points, an aliquot of the resulting mixture is injected into an HPLC to obtain the ratio of drug ester to drug acid or drug alcohol. An Arrhenius plot of the data provides a $t_{1/2}$ for hydrolysis. Table 2 below provides $t_{1/2}$ values for a variety of compounds.

Table 2

COMPOUND	$t_{1/2}$
Ketoprofen Methyl Ester	>48 h
Ketoprofen Ethyl Ester	>48 h
Ketoprofen Norcholine Ester	315 min.
Ketorolac Methyl Ester	>48 h
Ketorolac Ethyl Ester	>48 h
Ketorolac Norcholine Ester	14 min
Indomethacin Methyl Ester	>48 h
Indomethacin Ethyl Ester	>48 h
Indomethacin Norcholine Ester	315 min.
Apomorphine Diacetate	>48 h

10

EXAMPLE 7*General Procedure for Human Serum Hydrolysis Studies of Drug Esters*

Human serum (2.34 mL) is placed in a test tube. To the serum is added 260 μ L of a 10 mM solution of drug ester in acetonitrile. The tube is placed in a 37 °C incubator, and at various time points a 500 μ L aliquot is removed. The aliquot is mixed with 500 μ L methanol, and the mixture is vortex mixed and centrifuged. A sample of the supernatant is analyzed by HPLC obtain the ratio of drug ester to drug acid or drug alcohol. An

15

Arrhenius plot of the data provides a $t_{1/2}$ for hydrolysis. Table 3 below provides $t_{1/2}$ values for a variety of compounds.

Table 3

COMPOUND	$t_{1/2}$
Ketoprofen Methyl Ester	144 min
Ketoprofen Ethyl Ester	224 min
Ketoprofen Norcholine Ester	37 s
Ketorolac Ethyl Ester	90 min
Ketorolac Norcholine Ester	13 s
Indomethacin Methyl Ester	>48 h
Indomethacin Ethyl Ester	>48 h
Indomethacin Norcholine Ester	23 min
Apomorphine Diacetate	76.2 s

5

EXAMPLE 8

General Procedure for Screening Drug Esters for Aerosolization Preferability

Drug ester (1 mg) is dissolved or suspended in a minimal amount of a suitable solvent (e.g., dichloromethane or methanol). The solution or suspension is pipetted onto the middle portion of a 3 cm by 3 cm piece of aluminum foil. The coated foil is wrapped around the end of a 1½ cm diameter vial and secured with parafilm. A hot plate is preheated to approximately 300 °C, and the vial is placed on it foil side down. The vial is left on the hotplate for 10 s after volatilization or decomposition has begun. After removal from the hotplate, the vial is allowed to cool to room temperature. The foil is removed, and the vial is extracted with dichloromethane followed by saturated aqueous NaHCO₃. The organic and aqueous extracts are shaken together, separated, and the organic extract is dried over Na₂SO₄. An aliquot of the organic solution is removed and injected into a reverse-phase HPLC with detection by absorption of 225 nm light. A drug ester is preferred for aerosolization where the purity of the drug ester aerosol isolated by this method is greater

than 85%. Such a drug ester has a decomposition index less than 0.15. The decomposition index is arrived at by subtracting the percent purity (*i.e.*, 0.85) from 1.

CLAIMS

1. An aerosol for inhalation therapy, wherein the aerosol comprises particles comprising at least 10 percent by weight of a drug ester.

2. The aerosol according to Claim 1, wherein the drug ester is selected from a group of esters consisting of the following: esters of antibiotics; esters of anticonvulsants; esters of antidepressants; esters of antihistamines; esters of antiparkinsonian drugs; esters of drugs for migraine headaches; esters of drugs for the treatment of alcoholism; esters of muscle relaxants; esters of anxiolytics; esters of nonsteroidal anti-inflammatories; esters of other analgesics; and, esters of steroids.

3. The aerosol according to Claim 1, wherein the drug ester is an ester of a nonsteroidal anti-inflammatory.

4. The aerosol according to Claim 1, wherein the drug ester is an ester of an antiparkinsonian drug.

5. The aerosol according to Claim 3, wherein the nonsteroidal anti-inflammatory is selected from a group consisting of ketoprofen, ketorolac, and indomethacin.

6. The aerosol according to Claim 4, wherein the antiparkinsonian drug is apomorphine.

7. The aerosol according to Claim 5, wherein the ester is a methyl ester, an ethyl ester, or a norcholine ester.

8. The aerosol according to Claim 5, wherein the ester is apomorphine diacetate.

9. A method of delivering a drug ester to a mammal through an inhalation route, wherein the route comprises:

- a) heating a composition, wherein the composition comprises at least 5 percent by weight of drug ester, to form a vapor; and,
- b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles,

5 which is inhaled by the mammal.

10 10. The method according to Claim 9, wherein the particles comprise at least 10 percent by weight of drug ester.

10 11. The method according to Claim 9, wherein the drug ester is selected from a group of esters consisting of the following: esters of antibiotics; esters of anticonvulsants; esters of antidepressants; esters of antihistamines; esters of antiparkinsonian drugs; esters of drugs for migraine headaches; esters of drugs for the treatment of alcoholism; esters of muscle relaxants; esters of anxiolytics; esters of nonsteroidal anti-inflammatories; esters of
15 other analgesics; and, esters of steroids.

12. The method according to Claim 9, wherein the drug ester is an ester of a nonsteroidal anti-inflammatory.

20 13. The method according to Claim 9, wherein the drug ester is an ester of an antiparkinsonian drug.

14. The method according to Claim 12, wherein the nonsteroidal anti-inflammatory is selected from a group consisting of ketoprofen, ketorolac, and
25 indomethacin.

15. The method according to Claim 13, wherein the antiparkinsonian drug is apomorphine.

30 16. The method according to Claim 14, wherein the ester is a methyl ester, an ethyl ester, or a norcholine ester.

17. The method according to Claim 15, wherein the ester is apomorphine diacetate.

18. A kit for delivering a drug ester through an inhalation route to a mammal,
5 wherein the kit comprises:

- a) a composition comprising at least 5 percent by weight of drug ester; and,
- b) a device that forms a drug ester aerosol from the composition, for inhalation by the mammal

and wherein the device comprises:

- 10 a) an element for heating the drug ester composition to form a vapor;
- b) an element allowing the vapor to cool to form an aerosol; and,
- c) an element permitting the mammal to inhale the aerosol.

15




Fig. 1

INTERNATIONAL SEARCH REPORT

Int  al Application No

PCT/US 02/18543

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/72 A61K31/192 A61K31/407 A61K31/405 A61K31/473

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 607 691 A (OLDENBURG KEVIN R ET AL) 4 March 1997 (1997-03-04) cited in the application column 14, line 19 - line 28 column 33, line 20 - line 45 column 35, line 62 - column 36, line 26 column 53, line 11 - line 52 example 6 tables 3,5	1-3,5,7
A	---	4,6,8-18
X	EP 0 039 369 A (SCHERING CORP) 11 November 1981 (1981-11-11) page 6, line 9 - line 24 page 13	1,2
A	---	3-18

	-/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

21 November 2002

Date of mailing of the international search report

29/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Giménez Miralles, J

INTERNATIONAL SEARCH REPORT

Int. Patent Application No
PCT/US 02/18543

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 112 598 A (BIESALSKI HANS K) 12 May 1992 (1992-05-12) column 2, line 51 -column 5, line 14 examples	1,2
A	-----	3-18
X	US 5 126 123 A (JOHNSON KEITH A) 30 June 1992 (1992-06-30) column 3, line 34 -column 4, line 55 examples	1,2
A	-----	3-18
X	WO 00 28979 A (SKYEPHARMA AG ;MUELLER WALZ RUDI (DE); KELLER MANFRED (DE)) 25 May 2000 (2000-05-25) page 12, line 3 -page 14, line 13	1,2
A	-----	3-18

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 9-17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 9-17

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US 02/18543

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9-17
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/US 02/18543

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 5607691	A	04-03-1997	AU	6764794 A	29-08-1994
			CA	2153243 A1	18-08-1994
			EP	0683668 A1	29-11-1995
			JP	8510720 T	12-11-1996
			WO	9417792 A2	18-08-1994
			US	5814603 A	29-09-1998
			US	5622944 A	22-04-1997
			AU	4534593 A	04-01-1994
			EP	0647133 A1	12-04-1995
			WO	9325197 A1	23-12-1993
EP 0039369	A	11-11-1981	AU	533551 B2	01-12-1983
			EP	0039369 A1	11-11-1981
			AT	3774 T	15-06-1983
			AU	5810780 A	19-11-1981
			DE	3063750 D1	21-07-1983
US 5112598	A	12-05-1992	DE	3815221 A1	23-11-1989
			AT	99928 T	15-01-1994
			DE	58906676 D1	24-02-1994
			EP	0352412 A2	31-01-1990
			ES	2061772 T3	16-12-1994
			JP	2056419 A	26-02-1990
			JP	2766666 B2	18-06-1998
			US	5556611 A	17-09-1996
US 5126123	A	30-06-1992	AT	196076 T	15-09-2000
			AU	649702 B2	02-06-1994
			AU	8213591 A	23-01-1992
			CA	2083821 A1	29-12-1991
			DE	69132407 D1	12-10-2000
			DE	69132407 T2	05-04-2001
			EP	0536250 A1	14-04-1993
			ES	2151476 T3	01-01-2001
			GB	2263064 A ,B	14-07-1993
			IE	912246 A1	01-01-1992
			JP	3056784 B2	26-06-2000
			KR	175164 B1	01-02-1999
			NZ	238749 A	27-07-1993
			PT	98105 A ,B	31-08-1993
			RU	2098082 C1	10-12-1997
			WO	9200107 A1	09-01-1992
			ZA	9104957 A	29-04-1992
WO 0028979	A	25-05-2000	AU	6457899 A	05-06-2000
			WO	0028979 A1	25-05-2000
			CN	1326341 T	12-12-2001
			CZ	20011553 A3	12-09-2001
			EP	1131059 A1	12-09-2001
			HU	0104226 A2	28-02-2002
			JP	2002529498 T	10-09-2002
			NO	20012346 A	26-06-2001
			NZ	511527 A	25-10-2002
			PL	347640 A1	22-04-2002
			SK	6322001 A3	07-01-2002